

S. I. Kosenko, A. A. Kuznetsova, L. A. Vitavetskaya, Yu. G. Chernyakova, S. S. Seredenko

Odessa National Politechnical University, 1, Shevchenko av., Odessa, Ukraine

Odessa National Maritime Academy, Odessa, 4, Didrikhsona str., Odessa, Ukraine

Odessa State Environmental University, 15, Lvovskaya str., Odessa, Ukraine

e-mail: nucvita@mail.ru

CALCULATING THE RADIATIVE VACUUM POLARIZATION CONTRIBUTION TO THE ENERGY SHIFT OF SOME TRANSITIONS IN PIONIC AND KAONIC NITROGEN

Calculating the radiative contribution due to the vacuum polarization effect to energy value for some transitions in pionic and kaonic nitrogen has been carried out using the modified Uehling-Serber potential. The master equation for computing spectrum of the kaonic atoms is the Klein-Gordon-Fock equation. The vacuum polarization contribution to the 8-7 transition energy in kaonic nitrogen is calculated

This paper goes on our work on estimating the vacuum-polarization contributions to the transition energies of the hadronic atoms that is of a great importance especially in the last years (look below) [1-15]. Here we study pionic and kaonic nitrogen. Let us remind [13] that due to the significant progress in the modern experimental technologies now a great interest attracts studying spectra of heavy and super heavy elements atoms, exotic atomic systems, including hadronic and leptonic atoms [1-15]. Especial problem is connected with précised calculating the radiative corrections to the transition energies of the low- Z exotic (pionic, kaonic, muonic) atoms, namely, hydrogen and deuterium. Naturally, it is provided by necessity of further developing the modern as atomic and as nuclear theories. From the other side, detailed information about spectra of the exotic atomic systems (kaonic, pionic, muonic atoms) can be very useful under construction of the new X-ray standards. One could remind a great importance of the muonic chemistry, muonic spectroscopy. Very attractive perspective of the thermonuclear fission through the mechanism of the muonic catalysis is still interesting and widely studied.

The standard Dirac approach is traditionally used as starting basis in calculations of the heavy ions [2]. The problem of accounting the radiative corrections, in particular, self-energy part of the Lamb shift and vacuum polarization contribution is mostly treated with using the expansions on the natural physical parameters $1/Z$, αZ (α is fine structure constant) [5,10]. It permits evaluations of the relative contributions of different expansion energy terms: non-relativistic, relativistic ones, as functions of Z . For high Z (Z is a nuclear charge) it should be necessary to account for the high-order QED corrections and the nuclear finite size correction etc [1-3,10-12,16]. Further improvement of this method in a case of the heavy ions is linked with using gauge invariant procedures of generating relativistic orbital bases and more correct treating nuclear and QED effects [1-3]. In a case of the low- Z exotic atomic systems such as an exotic hydrogen (deuterium) a great interest attracts estimation of the radiative, in particular, vacuum polarization, correction. In refs. [17-19] it has been proposed a precise scheme to calculating spectra of heavy systems with account of nuclear and radiative effects, based on the relativistic many-body perturbation theory (see also

[3]) and advanced effective procedures for accounting the radiative corrections.

In this paper we present the results of calculating the contribution due to the vacuum polarization effect to energy shift for some transitions in pionic and kaonic nitrogen. The obtained result is compared with calculation data by Indelicato [19]. The details of the calculation procedure have been presented earlier, and here we consider only the key topics. The calculation of the radiative vacuum polarization shift in the pionic deuterium should be performed using the Dirac approximation as a zeroth one. Further, the expectation value of the radiative vacuum polarization operator gives the corresponding correction. The total electromagnetic interaction potential:

$$V(r) = V_n(r) + U(r). \quad (1)$$

includes the electrical V_n and polarization $U(r)$ potentials of a nucleus with accounting the finite size correction. As usually, the Coulomb potential for spherically symmetric density $\rho(r|R)$ can be written as follows:

$$V_n(r|R) = -\left(\frac{1}{r}\right) \int_0^r dr' r'^2 \rho(r'|R) + \int_r^\infty dr' r' \rho(r'|R). \quad (2)$$

The details of the determination of this potential can be, for example, found in ref. [21,22]. The vacuum polarization part is usually accounted in the first PT order by using the Uehling potential [1,8,16,17]:

$$U(r) = -\frac{2\alpha}{3\pi} \int_1^\infty dt \exp(-2rt/\alpha Z) \left(1 + 1/2t^2\right) \frac{\sqrt{t^2-1}}{t^2} \equiv$$

$$= -\frac{2\alpha}{3\pi} C(g) \quad (3)$$

$$g = \frac{r}{\alpha Z}. \quad (4)$$

The corresponding expectation value of this operator gives the corresponding vacuum polarization correction. In the scheme [12] this potential is approximated by quite precise analytical function (see details in refs. [3,16,17]). The most advanced version of the such potential ($C \rightarrow \tilde{C}$) is presented as follows:

$$\tilde{C}(g) = \tilde{C}_1(g) \tilde{C}_2(g) / \left(\tilde{C}_1(g) + \tilde{C}_2(g) \right),$$

$$\tilde{C}_2(g) = \tilde{C}_2(g) f(g),$$

$$\tilde{C}_2(g) = -1.8801 \exp(-g) / g^{3/2}$$

$$\tilde{C}_1(g) = h(g/2) + 1.410545 - 1.037837g,$$

$$f(g) = ((1.1024/g - 1.3361)/g + 0.8027)$$

The using this formula permits one to decrease the calculation errors for this term down to $\sim 0.1\%$. Error of usual calculation scheme is $\sim 10\%$.

We carried out the calculation of the vacuum polarization contribution to the energy shift for 8k-7 transition in kaonic nitrogen. It should be noted that the energy levels of exotic (kaonic) muonic atoms are very sensitive to effects of QED, nuclear structure and recoil since the kaon is heavier than the electron. As usually the fundamental constants from the CODATA 1998 are used in the numerical calculations.

We have evaluated the modified Uehling-Serber potential expectation values and obtained the value for the vacuum-polarization correction: 1.1783 eV. It is interesting to compare this result with the result by Indelicato, namely, the total vacuum polarization contribution: 1.1789 eV and from ref. [23]: 1,1778 eV. Further, let us present the data for the main Coulomb contribution from [8,24] - 2968.4565 eV and from ref. [23] - 2968.4492eV. Our estimate: 2968.4481eV. Further we have evaluated the modified Uehling-Serber potential expectation values and obtained the value for the vacuum-polarization correction into the 5f-4d transition energy in the pionic nitrogen. Our value for the vacuum-polarization correction: 2,9467eV. It is interesting to compare this result with the result by Serga [25] 3,216 eV. It should be noted that the last result is received with account of the Wichman-Kroll and Kallen-Sabry corrections. In any case, the physically reasonable agreement can be easily explained by the fact that pionic and kaonic nitrogen is the low-Z atomic system and in this case the expansion on the parameter αZ works sufficiently well.

References

1. Botham C., Martensson A.M., Sanders P.G. Relativistic effects in atoms and molecules.-Vancouver: Elseiver,2001.-550p.
2. Grant I.P., Relativistic Quantum Theory of Atoms and Molecules.-Oxford, 2008.-650P.
3. Glushkov A.V., Relativistic quantum theory. Quantum mechanics of atomic systems, Odessa: Astroprint, 2008.-700P.
4. Hayano R.S., Hori M., Horvath D., Widman E., Antiprotonic helium and CPT invariance//Rep. Prog. Phys.-2007.-Vol.70.-P.1995-2065.
5. Schlessler S., Le Bigot E.-O., Indelicato P., Pachucki K., Quantum-electrodynamics corrections in pionic hydrogen// Phys.Rev.C.-2011.-Vol.84.-P.015211 (8p.).
6. Gotta D., Amaro F., Anagnostopoulos D., Pionic Hydrogen//Precision Physics of Simple Atoms and Molecules, Ser. Lecture Notes in Physics (Springer, Berlin / Heidelberg).-2008.-Vol.745.-P.165-186.
7. Mitroy J., Quantum defect theory for the study of hadronic atoms/ Mitroy J., Ivallov I.A.// J. Phys. G: Nucl. Part. Phys.-2001.-Vol.27.-P.1421-1433
8. Deslattes R., Kessler E., Indelicato P., de Billy L., Lindroth E., Anton J., Exotic atoms//Rev. Mod. Phys. -2003.-Vol.75.-P.35-70.
9. Borie E., Lamb shift in muonic hydrogen// Phys.Rev. A.-2005.-Vol.71.-P.032508-1-8.
10. Mohr P.J. Quantum Electrodynamics Calculations in few-Electron Systems// Phys. Scripta.-1993.-Vol.46,N1.-P.44-52.
11. Klaft I.,Borneis S., Engel T., Fricke B., Grieser R., Huber G., Kuhl T., Marx D., Neumann R., Schroder S., Seelig P., Volker L. Precision laser spectroscopy of ground state hyperfine splitting of H-like $^{209}\text{Bi}^{82+}$ // Phys.Rev.Lett.-1994.-Vol.73.-P.2425-2427
12. Khetselius O.Yu., Relativistic perturbation theory calculation of the hyperfine structure parameters for some heavy-element isotopes// Int. Journ. of Quantum Chemistry.-2009.-Vol.109.- N14.-P. 3330-3335.
13. Kuznetsova A.A., Vitavetskaya L.A., Chernyakova Yu.G., Korchevsky D.A., Calculating radiative vacuum polarization contribution to the energy shift of 3p-1s transition in pionic deuterium// Photo-electronics.-2013.-Vol.22.-P.108-111
14. Backenstoss G., Pionic atoms//Ann.Rev. Nucl.Sci.-1990.-Vol.20.-P.467-510.
15. Menshikov L I and Evseev M K, Some questions of physics of exotic atoms// Phys. Uspekhi.2001.-Vol. 171.-P.150-184.
16. Ivanova E.P., Ivanov L.N., Aglitsky E.V., Modern trends in spectroscopy of multicharged ions// Phys.Rep.-1991.-Vol.166.-P.315-390.
17. Glushkov A.V., Ambrosov S., Loboda A., Chernyakova Y.G., Khetselius O., Svinarenko A., QED theory of the superheavy elements ions: energy levels, radiative corrections, and hyperfine structure for different nuclear models// Nuclear Phys.A-2004.-Vol. 734.-P.21-28.
18. Glushkov A.V., Khetselius O.Yu., Gurnitskaya E.P., Loboda A.V., Florko T.A., Sukharev D.E., Lovett L., Gauge-invariant QED perturbation theory approach to calculating nuclear electric quadrupole moments, hfs structure constants for heavy atoms and ions// Frontiers in Quantum Systems in Chem. and Phys. (Springer).-2008.-Vol.18.- P.505-522.
19. Vitavetskaya L.A., Malinovskaya S.V., Dubrovskaya Yu., Advanced quantum calculation of beta decay probabilities// Low Energy Antiproton Phys.-2005.-Vol.796.- P.201-205.
20. Gurnitskaya E.P., Khetselius O.Yu., Loboda A.V., Vitavetskaya L.A., Consistent quantum approach to quarkonium energy spectrum and semiconductor superatom and in external electric field// Photoelectronics.-2008.-N17.-P.127-130.

21. Kvasikova A.S., Ignatenko A.V., Florko T.A., Sukharev D.E., Chernyakova Yu.G., Photoeffect and spectroscopy of the hydrogen atom in the crossed dc electric and magnetic field// Photoelectronics.-2011.-Vol.20.-P.71-75.
22. Kuznetsova A.A., Kvasikova A.S., Shakhman A.N., Vitavetskaya L.A., Calculating the radiative vacuum polarization contribution to the energy shift of 2p-2s transition in μ -hydrogen// Photoelectronics.-2012.-N21.-P.66-67.
23. Khetselius O.Yu., Turin A.V., Sukharev D.E., Florko T.A., Estimating of X-ray spectra for kaonic atoms as tool for sensing the nuclear structure// Sensor Electr. and Microsyst. Techn.-2009.-N1.-P.30-35..
24. Indelicato P., Pionic nitrogen// Private communication.
25. Serga I.N., Relativistic theory of spectra of pionic atoms with account of the radiative and nuclear corrections// Photoelectronics.-2010 - Vol.22.-P.84-92

This article has been received within 2014

UDC 539.184

S. I. Kosenko, A. A. Kuznetsova, L. A. Vitavetskaya, Yu. G. Chernyakova, S. S. Seredenko

CALCULATING THE RADIATIVE VACUUM POLARIZATION CONTRIBUTION TO THE ENERGY SHIFT OF SOME TRANSITIONS IN PIONIC AND KAONIC NITROGEN

Abstract

Calculating the radiative contribution due to the vacuum polarization effect to energy value for some transitions in pionic and kaonic atoms has been carried out using the modified Uehling-Serber potential. The master equation for computing spectrum of the kaonic atoms is the Klein-Gordon-Fock equation. The vacuum polarization contribution to the 8-7 transition energy in kaonic nitrogen is calculated.

Key words: kaonic nitrogen, radiative corrections

УДК 539.184

С. И. Косенко, А. А. Кузнецова, Л. А. Витавецкая, Ю. Г. Чернякова, С. С. Середенко

РАСЧЕТ РАДИАЦИОННОГО ВКЛАДА ЗА СЧЕТ ЭФФЕКТА ПОЛЯРИЗАЦИИ ВАКУУМА В СДВИГ ЭНЕРГИИ РЯДА ПЕРЕХОДОВ В ПИОННОМ И КАОННОМ АЗОТЕ

Резюме

Проведен расчет радиационного вклада за счет эффекта поляризации вакуума в величину энергии ряда переходов в пионном и каонном атомах с использованием модифицированного потенциала Юлинга-Сербера. Базисным для расчета спектра каонных атомов является уравнение Клейна-Гордона-Фока. Рассчитан вклад за счет поляризации вакуума в энергию 8-7 перехода в каонном азоте.

Ключевые слова: каонный азот, радиационные поправки

С. І. Косенко, Г. О. Кузнецова, Л. А. Вітавецька, Ю. Г. Чернякова, С. С. Середенко

**РОЗРАХУНОК РАДІАЦІЙНОГО ВНЕСКУ ЗА РАХУНОК ЕФЕКТУ ПОЛЯРИЗАЦІЇ
ВАКУУМУ У ЗСУВ ЕНЕРГІЇ ДЕКОТРИХ ПЕРЕХОДІВ У ПІОННОМУ І КАОННОМУ
АЗОТЕ**

Резюме

Виконано розрахунок радіаційного внеску за рахунок ефекту поляризації вакууму у величину енергії декотрих переходів у піонному і каонному атомах з використанням модифікованого потенціалу Юлінга-Сербера . Базисним для розрахунку спектра каонних атомів є рівняння Клейна-Гордона-Фока. Розраховано внесок за рахунок поляризації вакууму в енергію 8-7 переходу у каонному азоті.

Ключові слова: каонний азот, радіаційні поправки