

International Research-to-Practice Conference
Climate Services: Science and Education

September 22-24, 2021
Odesa, Ukraine



Conference Proceedings



Co-funded by the
Erasmus+ Programme
of the European Union

DETECTION OF THE CLIMATE CHANGE IMPACT ON THE RIVER RUNOFF OF SPRING FLOOD IN PIVDENNY BUG RIVER BASIN

A.O. Dokus, PhD, Zh.R. Shakirzanova, Prof., Sc.D.

Odessa State Environmental University

In this work the climate change impact on the river runoff of spring flood is detected and the regional methodology of long-term forecasts of characteristics of spring flood of the rivers in the Pivdenny Buh river basin is developed. The methodology forecasts based on the preliminary typification of floods by water content in choosing the main factors of spring runoff formation, establishing regional parameters (within the basin zoning by a set of hydrometeorological factors) for long-term forecasting of runoff layers and maximum discharge, determining their probabilistic values over many years in climate change conditions.

Researches of temporal trends and cyclicity in long-term series of maximum runoff characteristics of spring floods in the Pivdenny Buh basin showed that the decrease of spring runoff of rivers in recent decades is caused by climate change due to rising air temperature in cold period, deficiency of stable snow covers and significant reducing the depth of soil freezing (since 1989). There is a rescheduling tendency of maximum snow reserves and spring floods to earlier dates (Dokus, Shakirzanova, 2017).

The time series of the maximum discharge and runoff layers of spring floods are synchronous in terms of cyclical fluctuations, and from the end of the 1980s to 2015 the low-water phase was observed on the rivers.

Analysis of the total curves of runoff layers and maximum discharge of spring flood and their difference integral curves in the Pivdenny Buh river basin (at posts with the largest operating hydro-electric power stations on them) showed that in the modern period of spring flood formation in the basin, the impact of river control on the maximum runoff is not significant, the deviations on the total curves are mainly explained by natural cycles and the impact of climate change on river water content.


In the research using the methods of factor and cluster analyzes; hydrological zoning was performed under the conditions of spring flood of rivers in the basin of the Pivdenny Buh, which allowed to distinguish two hydrological regions (region I and region II with subregions IIa, IIb, IIc). The analyzed natural conditions have shown that within limits of the carried out hydrological zoning these regions have specific features of spring floods formation. Data from long-term meteorological, agrometeorological and hydrological observations in the Pivdenny Buh river basin (as of 2020) were used as input information for the development of the methodology forecasts, using data from observations in the Velykyi Kuyalnyk and Tiligul rivers.

To assess the impact of future climate change on the water regime of rivers in the spring flood period, climate change coefficients were calculated (according to IPCC climate models (CORDEX project) of RCP 4.5 scenario for the period 2021-2050), which were taken into account in determining the basic

characteristics of forecasting methods – average long-term runoff layers and maximum spring discharge rates; and modeling of spring runoff layers in the Pivdenny Buh basin in affected period was made.

The results showed that in the Pivdenny Buh basin in the region I in the period 2021-2050 was expected to reduce the long-term average characteristics of runoff (runoff layers and maximum water flow) of spring floods on average by 23 % ($k_{ch}= 0.77$), and in region II: sub-district IIa – by 37 % ($k_{ch}= 0.63$), sub-region IIб – by 51 % ($k_{ch}= 0.49$), sub-region IIв – by 75 % ($k_{ch}= 0.25$).

The layers of spring water runoff modeled according to climatic models in the period of 2021-2050 were expected to be lower than their long-term average value (established taking into account), which was 23 mm (fig. 1). The formation of catastrophically high floods in the period 2021-2050 according to the regional methodology of long-term forecasts of characteristics of spring flood of the rivers in the Pivdenny Buh river basin is not predicted.



REFERENCES

1. Dokus A., Shakirzanova Zh. (2017). Forecasting the hydrological risks of floods using the method of forecasting the maximum runoff of spring high water Collected Papers «Geography in Global Context: Achievements and Challenges», Akaki Tsereteli State University Kutaisi, Georgia, 3-4 June 2017. P. 40-55.